



**REDES-AT and RAPAL Uruguay,**  
Address: San José 1423 / Maldonado 1858  
E-mail: prensa@redes.org.uy /  
rapaluy@chasque.net  
Tel. No: (05982) 9082730 - 4132989  
Fax No.: (05982) 4100985  
http://www.redes.uy  
http://www.chasque.net/rapaluy

*Prepared by Dioxin, PCBs and Waste Working Group of the International POPs Elimination Network (IPEN) Secretariat, REDES-AT (Uruguay), RAPAL (Uruguay) and Arnika Association (Czech Republic)*



## **Contamination of chicken eggs near the cement kilns in Minas, Uruguay by dioxins, PCBs and hexachlorobenzene**



# Contamination of chicken eggs near the cement kilns in Minas, Uruguay by dioxins, PCBs and hexachlorobenzene

## “Keep the Promise, Eliminate POPs!” Campaign Report

*Prepared by Dioxin, PCBs and Waste Working Group of the International POPs Elimination Network (IPEN) Secretariat, REDES-AT (Uruguay), RAPAL (Uruguay) and Arnika Association (Czech Republic)*

### Executive Summary

Free-range chicken eggs collected near two cement kilns in Minas, Uruguay showed levels of polychlorinated biphenyls (PCBs) expressed in WHO-TEQs that were two times higher than the proposed limit in European Union (EU). In addition, dioxin levels in the eggs exceeded the newly proposed EU action level and were almost two-fold higher than background levels.. To our knowledge, this study represents the first data about U-POPs in chicken eggs from Uruguay.

Cement kilns located close to the city of Minas were pointed out as potential sources of POPs releases in the region, but further monitoring of possible PCBs content in their fuels is needed. The companies apparently do not monitor the chlorine content of the materials they burn and do not measure the pollutants released during their operation. A stream that eventually serves as a source of drinking water runs near the plants. In 2004, a significant quantity of hyperthyroidism cases emerged in the area near the kilns along with other serious health problems. This study indicates that better monitoring of cement kiln operations as a potential U-POPs source is needed in developing countries and in countries with economies under transition in general.

The toxic substances measured in this study are slated for reduction and elimination by the Stockholm Convention which holds its first Conference of the Parties beginning 2 May 2005 in Uruguay. Uruguay is a Party to Convention since it ratified the Treaty in February 2004. The Convention mandates Parties to take specific actions aimed at eliminating these pollutants from the global environment. We view the Convention text as a promise to take the actions needed to protect Uruguayan and global public's health and environment from the injuries that are caused by persistent organic pollutants, a promise that was agreed by representatives of the global community: governments, interested stakeholders, and representatives of civil society. We call upon Uruguay's governmental representatives and all stakeholders to honor the integrity of the Convention text and keep the promise of reduction and elimination of POPs.

### Recommendations

- 1) Include PCBs in the UNEP Toolkit and elevate the importance of PCBs releases in the guidelines for Best Available Techniques (BAT) and Best Environmental Practices and all other documents prepared under the Stockholm Convention.
- 2) More POPs monitoring in Uruguay is needed;
- 3) More publicly accessible data about U-POPs releases from cement kilns in developing countries and countries with economies under transition are needed to address these sources of U-POPs properly;
- 4) A PCBs releases inventory would help to properly address all sources of their releases in Uruguay and may help identify sources useful to other Latin American countries;

5) Stringent limits for PCBs in both fuel and waste as well as air emissions should be introduced into both national legislation and under international treaties.

6) Prohibit cement kilns from firing hazardous waste including fuels containing POPs or chlorinated substance that could lead to POPs formation.

## **Introduction**

Persistent organic pollutants (POPs) harm human health and the environment. POPs are produced and released to the environment predominantly as a result of human activity. They are long lasting and can travel great distances on air and water currents. Some POPs are produced for use as pesticides, some for use as industrial chemicals, and others as unwanted byproducts of combustion or chemical processes that take place in the presence of chlorine compounds. Today, POPs are widely present as contaminants in the environment and food in all regions of the world. Humans everywhere carry a POPs body burden that contributes to disease and health problems.

The international community has responded to the POPs threat by adopting the Stockholm Convention in May 2001. The Convention entered into force in May 2004 and the first Conference of the Parties (COP1) will take place on 2 May 2005 in Uruguay. Uruguay ratified the Convention in February 2004.

The Stockholm Convention is intended to protect human health and the environment by reducing and eliminating POPs, starting with an initial list of twelve of the most notorious, the “dirty dozen.” Among this list of POPs there are four substances that are produced unintentionally (U-POPs): polychlorinated biphenyls (PCBs), hexachlorobenzene (HCB), polychlorinated dibenzo-p-dioxins (PCDDs) and dibenzofurans (PCDFs) The last two groups are simply known as dioxins.

The International POPs Elimination Network (IPEN) asked whether free-range chicken eggs might contain U-POPs if collected near potential sources of U-POPs named by the Stockholm Convention. The surroundings of two cement kilns near Minas, Uruguay were selected as a sampling site since cement kilns are known to be significant sources of U-POPs.<sup>1</sup> Chicken eggs were chosen for several reasons: they are a common food item; their fat content makes them appropriate for monitoring chemicals such as POPs that dissolve in fat; and eggs are a powerful symbol of new life. Free range hens can easily access and eat soil animals and therefore their eggs are a good tool for biomonitoring environmental contamination by U-POPs. This study is part of a global monitoring of egg samples for U-POPs conducted by IPEN and reflects the first data about POPs in eggs in Uruguay.

## **Materials and Methods**

Please see Annex 1.

## **Results and Discussion**

### **U-POPs in eggs sampled near Minas, Uruguay**

The results of the analysis of a pooled sample of 8 eggs collected within a 2 km distance from the cement kilns ANCAP and CUSCA (see further text about these plants) close to the city of Minas are summarized in Tables 1 and 2. Pooled sample fat content was measured at 10.7%.

The sampled eggs exceeded the newly proposed EU limit for PCBs by almost two-fold. In addition, the eggs exceeded the newly proposed EU limit for sum of PCDD/Fs and PCBs expressed in WHO-TEQ. The level of dioxins exceeded the proposed EU action level for these chemicals as well as the limit for dioxins content in egg fat for use as feedstuffs.

**Table 1: Measured levels of POPs in eggs collected near two cement kilns in Minas, Uruguay per gram of fat.**

	Measured level	Limits	Action level
PCDD/Fs in WHO-TEQ (pg/g)	2.18	3.0 <sup>a</sup>	2.0 <sup>b</sup>
PCBs in WHO-TEQ (pg/g)	3.75	2.0 <sup>b</sup>	1.5 <sup>b</sup>
Total WHO-TEQ (pg/g)	5.93	5.0 <sup>b</sup>	-
PCB (7 congeners) (ng/g)	29.00	200 <sup>c</sup>	-
HCB (ng/g)	1.40	200 <sup>d</sup>	-

Abbreviations: WHO, World Health Organization; TEQ, toxic equivalents; pg, picogram; g, gram; ng, nanogram.

<sup>a</sup> Limit set up in The European Union (EU) Council Regulation 2375/2001 established this threshold limit value for eggs and egg products. There is even more strict limit at level of 2.0 pg WHO-TEQ/g of fat for feedingstuff according to S.I. No. 363 of 2002 European Communities (Feedingstuffs) (Tolerances of Undesirable Substances and Products) (Amendment) Regulations, 2002.

<sup>b</sup> These proposed new limits are discussed in the document Presence of dioxins, furans and dioxin-like PCBs in food. SANCO/0072/2004.

<sup>c</sup> Limit used for example in the Czech Republic according to the law No. 53/2002 as well as in Poland and/or Turkey.

<sup>d</sup> EU limit according to Council Directive 86/363/EEC, level in brackets is proposed new general limit for pesticides residues (under which HCB is listed) according to the Proposal for a Regulation of the European Parliament and of the Council on maximum residue levels of pesticides in products of plant and animal origin, COM/2003/0117 final - COD 2003/0052.

Table 2 shows the levels of U-POPs in eggs expressed as fresh weight.

**Table 2: Measured levels of POPs in eggs collected near two cement kilns in Minas, Uruguay per gram of egg fresh weight.**

	Measured level	Limits	Action level
PCDD/Fs in WHO-TEQ (pg/g)	0.23	1 <sup>a</sup>	-
PCBs in WHO-TEQ (pg/g)	0.40	-	-
Total WHO-TEQ (pg/g)	0.64	-	-
PCBs (7 congeners) (ng/g)	3.10	-	-
HCB (ng/g)	0.15	-	-

<sup>a</sup> U.S. Department of Agriculture Food Safety and Inspection Service [Memo 8 July 1997] Advisory to Owners and Custodians of Poultry, Livestock and Eggs. Washington, DC:U.S. Department of Agriculture, 1997. FSIS advised in this memo meat, poultry and egg product producers that products containing dioxins at levels of 1.0 ppt in I-TEQs or greater were adulterated. There is even more strict EU limit at level of 0.75 pg WHO-TEQ/g of eggs fresh weight for feeding stuff according to S.I. No. 363 of 2002 European Communities (Feedingstuffs) (Tolerances of Undesirable Substances and Products) (Amendment) Regulations, 2002.

To our knowledge, the measurements of U-POPs in this study represent the first data on U-POPs in chicken eggs ever reported in Uruguay. The levels of dioxins exceeding the EU action level observed in the egg samples support the need for further monitoring and longer-term changes to eliminate chlorinated chemicals that serve as donors for PCBs, dioxins and furans releases in all environment compartments. As PCBs are not fully covered as U-POPs by all relevant documents and/or their drafts prepared under Stockholm Convention, it is very important to close this gap and introduce stricter rules for handling PCBs-containing wastes and fuels as well as for PCBs releases into the

environment. Finally, more data about U-POPs releases from cement kilns U-POPs in developing countries and countries with economies under transition are needed.

It is clear that among the U-POPs listed under the Stockholm Convention , PCBs are the main contaminants found in the eggs from Minas. HCB was almost at the background levels (see Annex 6).

### Comparison with other studies of eggs

We compared the levels of PCBs and PCDD/Fs measured in this study with data from other studies that also used pooled samples and/or expressed mean values of analyzed eggs (please see Annexes 2, 3 and 4). The data for eggs described in this report follow on the heels of a similar studies in Slovakia, Kenya, Czech Republic, Belarus, India (Uttar Pradesh), Tanzania, Senegal, Uruguay and Mexico released since 21 March 2005.<sup>2, 3, 4, 5, 6, 7, 8</sup> The dioxin levels in eggs in this study exceeded background levels by almost 2-fold (0.2 - 1.2 pg WHO-TEQ/g of fat).

Some other studies showing elevated levels of dioxins include samples near a chemical plant in Usti nad Labem, Czech Republic,<sup>9</sup> where comparable levels of dioxins (2.9 pg WHO-TEQ/g of fat) were found. Much higher levels (almost 15-times and more) were found near an old waste incinerator in Maincy, France<sup>10</sup> and an area affected by a spread mixture of waste incineration residues in Newcastle, UK,<sup>11</sup> with levels measured at 42.47 pg WHO-TEQ/g and 31 pg WHO-TEQ/g respectively.

PCBs levels expressed in WHO-TEQs in eggs collected near Minas are comparable to those found in free range eggs from neighborhood of Mbeubeuss dumpsite in Senegal<sup>12</sup> and/or from Beneshov in the Czech Republic (near a medical waste incinerator).

PCBs contribute over 60% of the whole WHO-TEQ value in these eggs as visible from the graph in Annex 5.

### Possible U-POPs sources

The elevated levels of PCBs and PCDD/Fs observed in this study provoke the question of possible sources. There are two major potential pollution sources: the two cement kilns. Another potential source of PCDD/Fs could also be open burning. Since there is no specific data about U-POPs measurements from the region, it is difficult to identify sources.

Since the major pollutant found in the eggs was PCBs, it is reasonable to question whether cement kilns can create PCBs. According to data from Poland shown in Table 3 they can. The data shows significant levels of PCBs produced from the Polish incinerator when burning coal or co-incinerating waste.

**Table 3: Measurements of POPs in cement kilns in Poland.**<sup>13</sup>

Cement kiln	PCDD/F ng-TEQ/m <sup>3</sup>	PCBs ng-TEQ/m <sup>3</sup>	HCB ng-TEQ/m <sup>3</sup>
Co-incinerating waste	0.070	8.95	44.2
Burning coal	0.055	4.45	2.90

One of the cement kilns near Minas in this study reportedly burns rice peel and fuel oil. Fuel oil can be sometimes contaminated by used oil with PCBs, but there is no data on the actual PCBs contents in these fuels. Monitoring data on the fuel and the cement kiln would help clarify these questions. In fact, more monitoring data on cements kilns in general is needed. Therefore we urge international as well as national institutions to strength U-POPs monitoring capacity in the region perhaps by developing a regional U-POPs monitoring project.

## **Minas and cement kilns**

### **Physical and administrative description, environmental and health consequences**

The potential pollution sources are two cement kilns located between 36 ° and 37 ° latitude south, and approx. 117 meters., altitude. The closest city to the cement kilns is “Minas”, which is 3-5 km away from from them at sea level., The winds can distribute the emissions from the kilns among the population in the urban and suburban area (see map at Picture 1 in Annex 1). Minas is the capitol of the department of “Lavalleja” in southeast Uruguay. It has a population of 37,149 habitants.

There is a stream. “La Plata” (The Silver), which passes only 20-50 meters from the cement plants. This stream runs down to another stream, “San Francisco”, which serves as a source of drinking water to the population of Minas and the adjacent area.

Several serious illnesses have been observed recently in Minas. In 2004, a significant quantity of cases of hyperthyroidism occurred in the neighbourhoods near the cement kiln. Strong evidence links hyperthyroidism to PBBs, PCBs, substituted phenols, dioxins, perchlorates, and the flame retardants, PBDEs.<sup>14</sup> According to the population living adjacent to the cement plants a series of health problems have been observed including: endocrine dysfunction, birth defects, abortions, respiratory and urinary problems. In addition, birth defects and abortions were also observed among the animals in the zone.

### **Cement kilns close to Minas in Uruguay**

Two different cement kilns are located within the studied area: 1) The ANCAP cement plant (Administración Nacional de Combustibles, Alcohol y Portland), a state owned company, located 5 km. from the urban zone of the city of Minas; and 2) The CUSCA cement plant (Compañía Uruguaya de Cemento Sociedad Anónima), a private company with investors from Spain.

The ANCAP, Pórtland cement factory located in Minas initiated operations in 1954 and according to to the compnay has two cement kilns with a production capacity 350 tons a day each. In 2003, this plant produced 120,000 tons of clinker. More data about ANCAP’s operation are in its annual reports.<sup>15</sup>

The ANCAP cement kiln burns fuel oil and rice peel. The combustion of rice peel does not generate solid residue and the fraction of inorganic matter is incorporated in the clinker. The combustion of these fuels in the kilns produces significant pollution: carbon dioxide, nitrogen oxide, sulphur dioxide and potentially others, but they are not measured.

The company admits that it does not regularly analyze the chlorine content of the rice peel or the fuel oil used as fuel. When a decision is made to use a cement kiln to burn POPs wastes or other halogenated wastes, the operator and national regulatory authorities should both be aware that this practice has the potential to generate and release large quantities of U-POPs to the environment as the Stockholm Convention correctly states.

The CUSCA cement kiln was installed in February 1997. It produces 1,400 tons of clinker daily and consumes 120 tons fuel oil per day.<sup>16</sup>

### **U-POPs and the Stockholm Convention**

The U-POPs measured in this study are slated for reduction and elimination by the Stockholm Convention which holds its first Conference of the Parties in May 2005 in Uruguay. Uruguay is a Party to Convention since it ratified the Treaty in 2004.

The Convention mandates Parties to take specific actions aimed at eliminating these pollutants from the global environment. Parties are to require the use of substitute or modified materials, products and processes to prevent the formation and release of U-POPs.<sup>a</sup> Parties are also required to promote the use of best available techniques (BAT) for new facilities or for substantially modified facilities in certain source categories (especially those identified in Part II of Annex C).<sup>b</sup> In addition, Parties are to promote both BAT and best environmental practices (BEP) for all new and existing significant source categories,<sup>c</sup> with special emphasis on those identified in Parts II and III. As part of its national implementation plan (NIP), each Party is required to prepare an inventory of its significant sources of U-POPs, including release estimates.<sup>d</sup> These NIP inventories will, in part, define activities for countries that will be eligible for international aid to implement their NIP. Therefore it is important that the inventory guidelines are accurate and not misleading.

The Stockholm Convention on POPs is historic. It is the first global, legally binding instrument whose aim is to protect human health and the environment by controlling production, use and disposal of toxic chemicals. We view the Convention text as a promise to take the actions needed to protect Uruguayan and global public's health and environment from the injuries that are caused by persistent organic pollutants, a promise that was agreed by representatives of the global community: governments, interested stakeholders, and representatives of civil society. We call upon Uruguayan governmental representatives and all stakeholders to honor the integrity of the Convention text and keep the promise of reduction and elimination of POPs.


---

<sup>a</sup> Article 5, paragraph (c)

<sup>b</sup> Article 5, paragraph (d)

<sup>c</sup> Article 5, paragraphs (d) & (e)

<sup>d</sup> Article 5, paragraph (a), subparagraph (i)



**Picture 1:** Map of Minas and surrounding with marked location of cement kilns (hot spot 1 = cement kiln ANCAP, hot spot 2 = cement kiln CUSCA and sampling places - 4 different chicken fanciers who provided their free range chicken eggs for IPEN's research.

## Annex 1. Materials and Methods

### Sampling

For sampling in Uruguay we have chosen the surroundings of two cement kilns close to the city of Minas.

The eggs were collected from four sites (see map at picture 1). The hens from which the eggs were picked were all free-range of age between 2 - 3 years although regularly provided with maize once at day, and the rest of their feeding is what they get from the soil.

Sampling was done by members of REDES-AT and RAPAL Uruguay on 20 January 2005. Four chicken fancier supplied 16 eggs from their free range chickens. The eggs were kept in cool conditions after sampling and then were boiled in Uruguay for 7 - 10 minutes in pure water and transported by express service to the laboratory at ambient temperature.


## **Analysis**

After being received by the laboratory, the eggs were kept frozen until analysis. The egg shells were removed and the edible contents of 6 eggs were homogenised. A 30 g sub-sample was dried with anhydrous sodium sulphate, spiked by internal standards and extracted by toluene in a Soxhlet apparatus. A small portion of the extract was used for gravimetric determination of fat. The remaining portion of the extract was cleaned on a silica gel column impregnated with H<sub>2</sub>SO<sub>4</sub>, NaOH and AgNO<sub>3</sub>. The extract was further purified and fractionated on an activated carbon column. The fraction containing PCDD/Fs, PCBs and HCB was analysed by HR GC-MS on Autospec Ultima NT.

Analysis for PCDD/Fs, PCBs and HCB was done in the Czech Republic in laboratory Axys Varilab. Laboratory Axys Varilab, which provided the analysis is certified laboratory by the Institute for technical normalization, metrology and probations under Ministry of Industry and Traffic of the Czech Republic for analysis of POPs in air emissions, environmental compartments, wastes, food and biological materials.<sup>a</sup> Its services are widely used by industry as well as by Czech governmental institutions. In 1999, this laboratory worked out the study about POPs levels in ambient air of the Czech Republic on request of the Ministry of the Environment of the Czech Republic including also soils and blood tests.


## Annex 2: Mean values found within different groups of eggs from different parts of world

Country/locality	Year	Group	Measured level in pg/g (WHO-TEQ) of fat	Source of information
3 EU countries (Ireland, Germany, Belgium)	1997-2003	both	0,63	DG SANCO 2004
Ireland, free range	2002-2005	free range	0,47	Pratt, I. et al. 2004, FSAI 2004
Ireland, organic eggs	2002-2005	free range	1,3	Pratt, I. et al. 2004, FSAI 2004
Belgium, Antwerp province	2004	free range	1,5	Pussemeier, L. et al. 2004
<b>Uruguay, Minas</b>	<b>2005</b>	<b>free range</b>	<b>2,18</b>	<b>Axys Varilab 2005</b>
Netherlands	2004	free range	2,6	SAFO 2004
UK, Newcastle	2002	free range	5,5	Pless-Mulloli, T. et al. 2003b
USA, Stockton	1994	free range	7,69	Harnly, M. E. et al. 2000
Belgium, Antwerp province, free range	2004	free range	9,9	Pussemeier, L. et al. 2004
Germany, Rheinfelden	1996	free range	12,7	Malisch, R. et al. 1996
USA, Oroville	1994	free range	18,46	Harnly, M. E. et al. 2000
France, Maincy	2004	free range	42,47	Pirard, C. et al. 2004
USA, Southern Mississippi, from grocery	1994	not free range	0,29	Fiedler, H. et al. 1997
Netherlands, commercial eggs	2004	not free range	0,3	Anonymus 2004
Ireland, barn eggs	2002-2005	not free range	0,31	Pratt, I. et al. 2004, FSAI 2004
Ireland, battery eggs	2002-2005	not free range	0,36	Pratt, I. et al. 2004, FSAI 2004
France, eggs from supermarkets	1995-99	not free range	0,46	SCOOP Task 2000
Sweden, commercial eggs	1995-99	not free range	1,03	SCOOP Task 2000
Germany, commercial eggs	1995-99	not free range	1,16	SCOOP Task 2000
Spain, supermarkets	1996	not free range	1,34	Domingo et al. 1999
Finland, commercial eggs	1990-94	not free range	1,55	SCOOP Task 2000
Belgium, Antwerp province, conventional farms	2004	not free range	1,75	Pussemeier, L. et al. 2004



### Annex 3: Levels of dioxins (PCDD/Fs) in different pool samples from different parts of world


Country/locality	Year	Group	Number of eggs/measured samples	Measured level in pg/g (WHO-TEQ) of fat	Source of information
UK, Newcastle (background level)	2000	free range	3/1 pooled	0,2	Pless-Mulloli, T. et al. 2001
Germany, Lower Saxony	1998	free range	60/6 pools	1,28	SCOOP Task 2000
UK, Newcastle (lowest level from pool samples)	2000	free range	3/1 pooled	1,5	Pless-Mulloli, T. et al. 2001
Uruguay, Minas	2005	free range		2,18	Axys Varilab 2005
Czech Republic, Usti nad Labem	2005	free range	6/1 pooled	2,9	Axys Varilab 2005
Tanzania, Vikuge	2005	free range	6/1 pooled	3,03	Axys Varilab 2005
Germany, Bavaria	1992	free range	370/37 pools	3,2	SCOOP Task 2000
Czech Republic, Klatovy	2003	free range	12	3,4	Beranek, M. et al. 2003
Belarus, Bolshoi Trostenec	2005	free range	6/1 pooled	3,91	Axys Varilab 2005
Czech Republic, Lysa nad Labem	2004	free range	4	6,8	Petrlik, J. 2005
Germany, Rheinfelden (lowest level from pool samples)	1996	free range	-	10,6	Malisch, R. et al. 1996
Slovakia, Kokshov-Baksha and Valaliky	2005	free range	6/1 pooled	11,52	Axys Varilab 2005
Germany, Rheinfelden (highest level from pool samples)	1996	free range	-	14,9	Malisch, R. et al. 1996
India, Lucknow	2005	free range	4/1 pooled	19,8	Axys Varilab 2005
Kenya, Dandora	2004	free range	6/1 pooled	22,92	Axys Varilab 2005
UK, Newcastle (highest level from pool samples)	2000	free range	3/1 pooled	31	Pless-Mulloli, T. et al. 2001
Senegal, Mbeubeuss	2005	free range		35,1	Axys Varilab 2005



## Annex 4: Levels of PCBs in WHO-TEQ in different chicken eggs samples from different parts of world


Country/locality	Year	Group	Number of measured samples	Specification	Measured level in pg/g (WHO-TEQ) of fat	Source of information
Netherlands, commercial eggs	1999	not free range	100/2 pools	pool, nonortho-PCBs	0,44	SCOOP Task 2000
Netherlands, organic farms (lowest level)	2002	free range	6	pool	0,7	Traag, W. et al. 2002
Tanzania, Vikuge	2005	free range	6/1 pool	pool	0,7	Axys Varilab 2005
Czech Republic, Klatovy-Luby	2003	free range	free range	individual	0,7	Beranek, M. et al. 2003
UK, commercial eggs	1992	not free range	24/1 pool	pool	0,97	SCOOP Task 2000
Czech Republic, Usti nad Labem	2005	free range	6/1 pool	pool	1,2	Axys Varilab 2005
Sweden, commercial eggs	1999	not free range	32/4 pools	pool	1,45	SCOOP Task 2000
Netherlands	1990	mixed	8/2 pools	pool, nonortho-PCBs	1,8	SCOOP Task 2000
Sweden, different eggs	1993	mixed	84/7 pools	pool	1,82	SCOOP Task 2000
UK, commercial eggs	1982	not free range	24/1 pool	pool	2,36	SCOOP Task 2000
Senegal, Mbeubeuss	2005	free range		pool	3,4	Axys Varilab 2005
Uruguay, Minas	2005	free range		pool	3,8	Axys Varilab 2005
Czech Republic, Beneshov	2004	free range	4	pool	3,9	Axys Varilab 2004
Uzbekistan, Kanlikul	2001	free range	-	individual	4,5	Muntean, N. et al. 2003
Slovakia, Kokshov-Baksha + Valaliky	2005	free range	6/1 pool	pool	4,6	Axys Varilab 2005
Netherlands, organic farms (highest level)	2002	free range	6	pool	5,76	Traag, W. et al. 2002
Kenya, Dandora	2004	free range	6/1 pool	pool	8,1	Axys Varilab 2005
India, Lucknow	2005	free range	4/1 pooled	pool	9,4	Axys Varilab 2005
Belarus, Bolshoi Trostenech	2005	free range	6/1 pool	pool	9,8	Axys Varilab 2005
Czech Republic, Lysa nad Labem	2004	free range	4	pool	22,4	Petrlik, J. 2005

## PCBs in WHO-TEQ



## Annex 5: Balance between PCDD/Fs versus PCBs in different eggs samples in WHO-TEQs


Country/locality	Year	Group	PCDD/Fs	PCBs	Total WHO-TEQ	Source of information
Czech Republic, Lysa nad Labem	2004	free range	6,80	22,40	29,20	Petrlik, J. 2005
Netherlands	2002	free range	4,74	5,76	10,50	Traag, W. et al. 2002
Netherlands	2002	free range	0,70	4,89	5,59	Traag, W. et al. 2002
Sweden	1993	mixed	1,31	1,82	3,13	SCOOP Task 2000
UK	1982	not free range	8,25	2,36	10,61	SCOOP Task 2000
UK	1992	not free range	1,77	0,97	2,74	SCOOP Task 2000
Sweden	1999	not free range	1,43	1,45	2,48	SCOOP Task 2000
Slovakia, Kokshov-Baksha + Valaliky	2005	free range	11,52	4,60	16,12	Axys Varilab 2005
Czech Republic, Usti nad Labem	2005	free range	2,9	1,22	4,12	Axys Varilab 2005
Tanzania, Vikuge	2005	free range	3,03	0,7	3,73	Axys Varilab 2005
Belarus, Bolshoi Trostenech	2005	free range	3,91	9,83	13,74	Axys Varilab 2005
India, Lucknow	2005	free range	19,8	9,4	29,2	Axys Varilab 2005
Senegal, Mbeubeuss	2005	free range	35,1	3,44	38,54	Axys Varilab 2005
Uruguay, Minas	2005	free range	2,18	3,75	5,93	Axys Varilab 2005



## Annex 6: Levels of HCB in ng/g of fat in different chicken eggs samples from different parts of world

Country	Date/year	Specification	Number of measured samples	Measured level in ng/g of fat	Source of information
Uzbekistan, Nukus	2001	free range	-		1,0Muntean, N. et al. 2003
Uruguay, Minas	2005	free range			1,4Axys Varilab 2005
Senegal, Mbeubeuss	2005	free range			1,7Axys Varilab 2005
India, Lucknow	2005	free range	4/1 pooled		3,8Axys Varilab 2005
Kenya, Dandora	2004	free range	6/1 pool		4,4Axys Varilab 2005
Belarus, Bolshoi Trostenech	2005	free range	6/1 pool		4,7Axys Varilab 2005
Slovakia, Kokshov-Baksha	2005	free range	6/1 pool		10,7Axys Varilab 2005
Czech Republic, Beneshov	2004	free range	4/1 pool		14,9Axys Varilab 2004
Slovakia, Stropkov, free range eggs	before 1999	free range	1		16,6Kocan, A. et al. 1999
Uzbekistan, Chimbay	2001	free range	-		19,0Muntean, N. et al. 2003
Tanzania, Vikuge	2005	free range	6/1 pool		19,1Axys Varilab 2005
Czech Republic, Usti nad Labem	2005	free range	6/1 pool		35,8Axys Varilab 2005
Slovakia, Michalovce, free range eggs	before 1999	free range	1		40,7Kocan, A. et al. 1999
Slovakia, Michalovce, commercial eggs	before 1999	not free range	1		2,7Kocan, A. et al. 1999
Slovakia, Stropkov, commercial eggs	before 1999	not free range	1		3,0Kocan, A. et al. 1999

## HCB in ng/g of fat



**Annex 7: Photos**



**Picture 1:** ANCAP Cement plant



**Picture 2:** CUCSA Cement plant



**Picture 3:** Sampling place 1.



**Picture 4:** Sampling place 2.



**Picture 5:** Sampling place 3.

## References

- 
- <sup>1</sup> Grochowalski, A. 2002: Sprawozdanie z przeprowadzonych pomiarów i oznaczania stężenia PCDDs/PCDFs, HCB i PCBs, 30.09.2002; <http://ks.ios.edu.pl/gef/doc/GF-POL-INV-R1.PDF>
- <sup>2</sup> Friends of the Earth, Arnika, IPEN Dioxin, PCBs and Waste WG 2005. Contamination of chicken eggs near the Koshice municipal waste incinerator in Slovakia by dioxins, PCBs, and hexachlorobenzene. Available at [www.ipen.org](http://www.ipen.org) 21 March 2005
- <sup>3</sup> ENVILEAD, Arnika, IPEN Dioxin, PCBs and Waste WG 2005. Contamination of chicken eggs near the Dandora dumpsite in Kenya by dioxins, PCBs, and hexachlorobenzene. Available at [www.ipen.org](http://www.ipen.org) 24 March 2005.
- <sup>4</sup> Arnika, IPEN Dioxin, PCBs and Waste WG 2005. Contamination of chicken eggs near the Spolchemie factory in Usti nad Labem in the Czech Republic by dioxins, PCBs, and hexachlorobenzene. Available at [www.ipen.org](http://www.ipen.org) 25 March 2005.
- <sup>5</sup> Foundation for Realization of Ideas, Arnika, IPEN Dioxin, PCBs and Waste WG 2005. Contamination of chicken eggs near the Bolshoy Trostenec dumpsite in Belarus by dioxins, PCBs, and hexachlorobenzene. Available at [www.ipen.org](http://www.ipen.org) 29 March 2005.
- <sup>6</sup> Toxics Link, Arnika, IPEN Dioxin, PCBs and Waste WG 2005: Contamination of chicken eggs near the Queen Mary's Hospital, Lucknow medical waste incinerator in Uttar Pradesh (India) by dioxins, PCBs, and hexachlorobenzene. Available at [www.ipen.org](http://www.ipen.org) 30 March 2005.
- <sup>7</sup> AGENDA, Arnika, IPEN Dioxin, PCBs and Waste WG 2005: Contamination of chicken eggs near the Vikuge obsolete pesticides stockpile in Tanzania by dioxins, PCBs, and hexachlorobenzene. Available at [www.ipen.org](http://www.ipen.org) 31 March 2005.
- <sup>8</sup> PAN Africa, Arnika, IPEN Dioxin, PCBs and Waste WG 2005: Contamination of chicken eggs near the Mbeubeuss dumpsite in a suburb of Dakar, Senegal by dioxins, PCBs and hexachlorobenzene. Available at [www.ipen.org](http://www.ipen.org) 4 April 2005.
- <sup>9</sup> Arnika, IPEN Dioxin, PCBs and Waste WG 2005. Contamination of chicken eggs near the Spolchemie factory in Usti nad Labem in the Czech Republic by dioxins, PCBs, and hexachlorobenzene. Available at [www.ipen.org](http://www.ipen.org) 25 March 2005.
- <sup>10</sup> Pirard, C., Focant, J.-F., Massart, A.-C., De Pauw, E., 2004: Assessment of the impact of an old MSWI. Part 1: Level of PCDD/Fs and PCBs in surrounding soils and eggs. Organohalogen Compounds 66: 2085-2090.
- <sup>11</sup> Pless-Mulloli, T., Edwards, R., Schilling, B., Paepke, O. 2001: Executive Summary. PCCD/PCDF and Heavy Metals in Soil and Egg Samples from Newcastle Allotments: Assessment of the role of ash from the Byker incinerator. (Includes comments from Food Standards Agency, Environment Agency). 12 February 2001. University of Newcastle.
- <sup>12</sup> PAN Africa, Arnika, IPEN Dioxin, PCBs and Waste WG 2005: Contamination of chicken eggs near the Mbeubeuss dumpsite in a suburb of Dakar, Senegal by dioxins, PCBs and hexachlorobenzene. Available at [www.ipen.org](http://www.ipen.org) 4 April 2005.

---

<sup>13</sup> Grochowalski, A. 2002: Sprawozdanie z przeprowadzonych pomiarów i oznaczania stężenia PCDDs/PCDFs, HCB i PCBs, 30.09.2002; <http://ks.ios.edu.pl/gef/doc/GF-POL-INV-R1.PDF>

<sup>14</sup> Janssen S, Solomon G, Schettler T. 2004, Chemical contaminants and human disease: A summary of the evidence. Natural Resources Defense Council and Science and Environmental Health Network. Available here: <http://www.protectingourhealth.org/corethemes/links/2004-0203spreadsheet.htm>

<sup>15</sup> ANCAP 2002: ANCAP - memoria y balance 2002. [http://www.ancap.com.uy/Balance/Mem\\_ANCAP2002.pdf](http://www.ancap.com.uy/Balance/Mem_ANCAP2002.pdf).

<sup>16</sup> Data provided by a local environmental NGO Arequita from Minas in January 2005.

---

## References for Tables in Annexes

Anonymus 2004: Analytical results eggs from both free range chickens and not free range chickens from Netherlands. Information provided by Netherlands to other EU member states. November 2004.

Axys Varilab CZ 2004: Protokoly č. 537/1-4 o stanovení PCDD/F, PCB vyjádřených ve WHO-TEQ, kongenerových PCB a HCB vydané zkušební laboratoří firmy Axys Vailab. Protocols No. 537/1-4. Vrane nad Vltavou, 2004.

Beranek, M., Havel, M., Petřík, J. 2003: Lindane - pesticide for the black list. Czech Ecological Society Report, Prague, Nov 2003.

DG SANCO 2004: Analysis of the data contained in the report "Dioxins and PCBs in Food and Feed : Data available to DG SANCO - Joint Report DG SANCO/DG-JRC-IRMM in the light of the proposed maximum levels in document SANCO/0072/2004.

Domingo, J.L., Schuhmacher, M., Granero, S., Llobet, J.M. 1999: PCDDs and PCDFs in food samples from Catalonia, Spain. An assessment of dietary intake. Chemosphere. 38(15):3517-3528. In US EPA 2000.

Fiedler, H.; Cooper, K.R.; Bergek, S.; Hjelt, M.; Rappe, C. (1997) Polychlorinated dibenzo-p-dioxins and polychlorinated dibenzofurans (PCDD/PCDF) in food samples collected in southern Mississippi, USA. Chemosphere. 34:1411-1419. In US EPA 2000.

Harnly, M. E., Petreas, M. X., Flattery, J., Goldman, L. R. 2000: Polychlorinated Dibenzo-p-dioxin and Polychlorinated Dibenzofuran Contamination in Soil and Home-Produced Chicken Eggs Near Pentachlorophenol Sources. Environ. Sci. Technol.2000, 34,1143-1149

Kočan, A., Jursa, S., Petřík, J., Drobná, B., Chovancová, J., Suchánek, P. 1999: Stav kontaminácie potravín polychlórovanými bifenylmi v zaťaženej oblasti okresu Michalovce a porovnávacej oblasti okresu Stropkov. In: Cudzorodé látky v potravinách, 10. - 12. máj 1999, Tatranská Štrba, pp. 31 - 32.

Malisch, R. 1998: Update of PCDD/PCDF-intake from food in Germany. Chemosphere. 37 (9 - 12):1687-1698. In US EPA 2000.

Malisch, R., Schmid, P., Frommberger, R., Fuerst, P. 1996: Results of a Quality Control Study of Different Analytical Methods for Determination of PCDD/PCDF in Eggs Samples. Chemosphere Vol. 32, No. 1, pp. 31-44.

---

Muntean, N., Jermini, M., Small, I., Falzon, D., Peter Fuerst, P., Migliorati, G., Scortichini, G., Forti, A. F., Anklam, E., von Holst, C., Niyazmatov, B., Bahkridinov, S., Aertgeerts, R., Bertollini, R., Tirado, C., Kolb, A. 2003: Assessment of Dietary Exposure to Some Persistent Organic Pollutants in the Republic of Karakalpakstan of Uzbekistan. Vol. 111, No 10, August 2003, Environmental Health Perspectives, 1306-1311.

Petrlík, J. 2005: Hazardous waste incinerator in Lysá nad Labem and POPs waste stockpile in Milovice. International POPs Elimination Project (IPEP) Hot Spot Report. Arnika, Prague 2005.

Pirard, C., Focant, J.-F., Massart, A.-C., De Pauw, E., 2004: Assessment of the impact of an old MSWI. Part 1: Level of PCDD/Fs and PCBs in surrounding soils and eggs. Organohalogen Compounds 66: 2085-2090.

Pless-Mulloli, T., Edwards, R., Schilling, B., Paepke, O. 2001b: Executive Summary. PCDD/PCDF and Heavy Metals in Soil and Egg Samples from Newcastle Allotments: Assessment of the role of ash from the Byker incinerator. (Includes comments from Food Standards Agency, Environment Agency). 12 February 2001. University of Newcastle.

Pless-Mulloli, T., Air, V., Schilling, B., Paepke, O., Foster, K. 2003b: Follow-up Assessment of PCDD/F in Eggs from Newcastle Allotments. University of Newcastle, Ergo, Newcastle City Council, July 2003.

Pratt, I., Tlustos, Ch., Moylan, R., Neilan, R., White, S., Fernandes, A., Rose, M. 2004: Investigation into levels of dioxins, furans and PCBs in battery, free range, barn and organic eggs. Organohalogen Compounds – Volume 66 (2004) 1925-31.

Pussemier, L., Mohimont, L., Huyghebaert, A., Goeyens, L., 2004. Enhanced levels of dioxins in eggs from free range hens: a fast evaluation approach. Talenta 63: 1273-1276.

SAFO (Sustaining Animal Health and Food Safety in Organic Farming) 2004: Onderzoek naar dioxine in eieren van leghennen met vrije uitloop. SAFO, September 2004. Published at: <http://www.agriholland.nl/nieuws/home.html>. 12/10/2004.

SCOOP Task 2000: Assessment of dietary intake of dioxins and related PCBs by the population of EU Member States. Reports on tasks for scientific cooperation Report of experts participating in Task 3.2.5 (7 June 2000) and Annexes to Report SCOOP Task 3.2.5 (Dioxins). Final Report, 7 June, 2000. European Commission, Health & Consumer Protection Directorate-General, Brussels 2000.

SVA CR (State Veterinary Administration of the Czech Republic) 2004: Chart with results of regular monitoring in Middle Bohemian region. Document reached by Arnika upon request for information.

Traag, W., Portier, L., Bovee, T., van der Weg, G., Onstenk, C., Elghouch, N., Coors, R., v.d. Kraats, C., Hoogenboom, R. 2002: Residues of Dioxins and Coplanar PCBs in Eggs of Free Range Chickens. Organohalogen Compounds Vol. 57 (2002). 245-248.